



**Experimental Mathematics** 

ISSN: 1058-6458 (Print) 1944-950X (Online) Journal homepage: www.tandfonline.com/journals/uexm20

# Gaps Between Consecutive Primes and the **Exponential Distribution**

Joel E. Cohen

To cite this article: Joel E. Cohen (2025) Gaps Between Consecutive Primes and the Exponential Distribution, Experimental Mathematics, 34:2, 260-269, DOI: 10.1080/10586458.2024.2362348

To link to this article: https://doi.org/10.1080/10586458.2024.2362348



Published online: 22 Jun 2024.



🕼 Submit your article to this journal 🕝



View related articles



View Crossmark data 🗹



#### Joel E. Cohen 💿

The Rockefeller University, New York, NY, USA; Departments of Statistics, Columbia University, New York, NY, USA; University of Chicago, Chicago, IL, USA

#### ABSTRACT

Based on the primes less than  $4 \times 10^{18}$ , Oliveira e Silva et al. (*Math. Comp.*, 83(288):2033–2060, 2014) conjectured an asymptotic formula for the sum of the *k*th power of the gaps between consecutive primes less than a large number *x*. We show that the conjecture of Oliveira e Silva holds if and only if the *k*th moment of the first *n* gaps is asymptotic to the *k*th moment of an exponential distribution with mean log *n*, though the distribution of gaps is not exponential. Asymptotically exponential moments imply that the gaps asymptotically obey Taylor's law of fluctuation scaling: variance of the first *n* gaps  $\sim$  (mean of the first *n* gaps)<sup>2</sup>. If the distribution of the first *n* gaps is asymptotic to  $(\log n)^2$ . The largest of the first *n* gaps is asymptotic to  $(\log n)^2$  if and only if the Cramér-Shanks conjecture holds. Numerical counts of gaps and the maximal gap  $G_n$  among the first *n* gaps test these results. While most values of  $G_n$  are better approximated by  $(\log n)^2$  than by other models, seven exceptional values of *n* with  $G_n > 2e^{-\gamma} (\log n)^2$  suggest that  $\limsup_{n\to\infty} G_n/[2e^{-\gamma} (\log n)^2]$  may exceed 1.

#### **KEYWORDS**

Cramér-Shanks conjecture; Firoozbakht's conjecture; fluctuation scaling; gap between consecutive primes; largest prime gap; Maier's theorem; prime gap; power variance function; Taylor's law; variance function

Taylor & Francis

Check for updates

Taylor & Francis Group

2020 MATHEMATICS SUBJECT CLASSIFICATION: Primary: 11A41; 11N05

#### 1. Introduction

The gaps between consecutive prime numbers have been studied mathematically, numerically, and statistically by many people for centuries [4, 8–10, 13, 16, 17, 19–21, 23–25, 28, 29, 32, 38–42]. This list of references is far from complete. Many questions remain open. Throughout, we use *gap* to mean a difference, denoted *d*, between *consecutive* primes, including  $d_4 := p_5 - p_4 = 11 - 7 = 4$  and excluding  $p_4 - p_2 = 7 - 3 = 4$ , where  $p_n$  is the *n*th prime,  $p_1 = 2$ .

We do not attempt to summarize here everything that is currently known about the distribution of prime gaps. Some selected recent results and conjectures are pertinent. Heath-Brown [21, p. 87] conjectured a simple asymptotic expression for the sum of the squares of the gaps of the primes not exceeding a large real *x* (OEIS <u>A074741</u>). Wolf [39, Conjecture 5] conjectured an alternative asymptotic form and compared the two alternatives numerically. Based on a detailed numerical study of the primes less than  $4 \times 10^{18}$ , Oliveira e Silva [28, p. 2056] conjectured a simple asymptotic formula for the sum of the *k*th power of the gaps of primes less than  $x \text{ as } x \to \infty$ , for every  $k = 1, 2, \ldots$ . Wolf [41] refined and extended the conjectured asymptotic expressions of Oliveira e Silva and compared the alternatives analytically and numerically. Firoozbakht conjectured that the sequence  $(p_n^{1/n})_{n \in \mathbb{N}}$  decreases as *n* increases [38]. The conjecture holds for all primes less than  $2^{64} \approx 1.84 \times 10^{19}$  as of April 2024 [38]. Kourbatov [23] showed that Firoozbakht's conjecture implies an upper bound on all gaps. Maier [24] showed that chains of consecutive large gaps exist, and that average gaps of primes in short intervals do not converge (contrary to previous conjecture) but have a limsup strictly greater than the liminf [25].

What is new here? We consider the first n = 1, 2, ... gaps instead of the gaps of primes less than large x (Section 3). This change of scale enables us to show (Theorem 3) that the conjecture of Oliveira e Silva holds if and only if, for every k = 1, 2, ..., the kth moment of the gaps (that is, the sum of the kth power of the first n gaps, divided by n) is asymptotically (as  $n \to \infty$ ) the kth moment  $k!(\log n)^k$  of an exponential distribution with mean  $\log n$ .

Obviously the distribution of gaps is not exponential because an exponential random variable has a continuous density function on  $[0, \infty)$ , whereas gaps are positive integers only and all gaps except  $d_1 = p_2 - p_1 = 3 - 2 = 1$  are even integers. Gaps are also not geometrically distributed (the integer-valued analog of the exponential) because a geometric distribution would give positive probability to all odd positive integers. Despite these deviations from the exponential distribution at a microscopic scale, the numerically supported conjecture that the *k*th moment of the first *n* gaps is asymptotically the *k*th moment  $k!(\log n)^k$  of an exponential distribution with mean log *n* has interesting consequences, as we now summarize and then show in detail.

From Theorem 3, it follows (Corollary 4) that, as  $n \to \infty$ , the ratio of the variance of the first *n* gaps to the square of the mean of the first *n* gaps converges to 1 (Section 4). Thus the gaps have a power-law asymptotic variance function [1, 2, 11, 34–36] or, equivalently, asymptotically obey Taylor's law of fluctuation scaling: variance of the first *n* gaps  $\sim$  square of the mean of the first *n* 

gaps [4, 6, 7, 12, 14, 33]. Moreover (Section 6), Taylor's law (with a different coefficient and a different exponent) also describes the variance function of the maximal gap among the first *n* gaps if the maximal gap may be modeled as the largest order statistic of *n* observations of an exponential distribution with mean log *n*: variance of the maximal order statistic  $\sim (\pi^2/6) \times$  mean of the maximal order statistic. These (conjectured) scaling laws do not appear to have been noticed previously.

The Cramér-Shanks conjecture (Cramér [8, pp. 24, 27], Shanks [32, p. 648, his Eq. (5)], Granville [17, p. 21, his Eq. (14)]) proposes that the maximal gap of primes less than x is asymptotic to  $(\log x)^2$  as  $x \to \infty$  (Section 6). Using the exponential distribution as a heuristic model of the gaps, we show (Theorem 5) that the Cramér-Shanks conjecture holds if and only if the largest of the first n gaps is asymptotic (as  $n \to \infty$ ) to the expectation  $(\log n)^2$  of the largest order statistic of a sample of n independent observations from an exponential distribution with mean  $\log n$ . However, the (1 - 1/n)-quantile of the largest order statistic of a sample of n independent observations from an exponential distribution with mean  $\log n$  yields  $2(\log n)^2$  as an asymptotic estimate of the largest of the first n gaps. This estimate exceeds the Cramér-Shanks conjecture and Kourbatov's bound [23] on gaps derived from Firoozbakht's conjecture.

In the concluding Section 7, we compare asymptotic expressions (derived from the exponential model of prime gaps and from other conjectures) for moments and maximal gaps with exact numerical results derived from public sources. Numerical evidence supports the moments of an exponential distribution with mean log *n* as models of the moments of the first *n* gaps and questions  $(\log n)^2$  as an asymptotic description of the largest of the first *n* gaps.

#### 2. Definitions and background

For any natural number  $n \in \mathbb{N} := \{1, 2, ...\}$ , let  $p_n$  be the *n*th prime starting from  $p_1 = 2$ ,  $p_2 = 3$  (OEIS A000040). Let  $\mathbb{P} := \{p_1, p_2, ...\}$  be the set of primes. Define the *n*th gap  $d_n$  between consecutive primes (OEIS A001223) and the sum  $D_k(x)$  of the *k*th power,  $k \in \{0\} \cup \mathbb{N}$ , of the gaps between consecutive primes that do not exceed real *x* to be

$$d_n := p_{n+1} - p_n, \ n \in \mathbb{N},\tag{1}$$

$$D_k(x) := \sum_{\substack{D_{n+1} \le x}} d_n^k.$$
<sup>(2)</sup>

We shall say that  $d_n$  is the gap of  $p_n$  if and only if (1). For example, the gap of 17 is 2.

Let  $\pi(x) := \#\{p \in \mathbb{P} \mid p \leq x\}$  be the number of primes that do not exceed real  $x \geq 2$  (OEIS <u>A000720</u> gives  $\pi(x)$  evaluated at positive integral values of x). Obviously  $\pi(p_n) = n$  and  $p_{\pi(x)} = \max\{p_n \in \mathbb{P} \mid p_n \leq x\}$ . For 2 < a < b, the number of primes in the interval (a, b] equals  $\pi(b) - \pi(a)$ . Those primes are  $p_{\pi(a)+1}, \ldots, p_{\pi(b)}$  and the gaps of those primes are  $d_{\pi(a)+1}, \ldots, d_{\pi(b)}$ . Then

$$d_{\pi(a)+1} + \dots + d_{\pi(b)-1} := L_1 < L := b - a < L_2 := d_{\pi(a)+1} + \dots + d_{\pi(b)}.$$
(3)

Some authors define the number of the gaps of the primes in (a, b] to be  $N_1 := \pi(b) - \pi(a) - 1$  (excluding gap  $d_{\pi(b)}$  because  $p_{\pi(b)+1} > b$ ) and some authors define it to be  $N_2 := \pi(b) - \pi(a)$  (including gap  $d_{\pi(b)}$ ). Asymptotically, the difference in these definitions is immaterial.

If f(x) and g(x) are real-valued functions of real x and g(x) > 0 for all x sufficiently large, define  $f(x) \sim g(x)$  to mean that  $f(x)/g(x) \rightarrow 1$  as  $x \rightarrow \infty$ . In Riesel's [29, p. 61] cautious notation, define  $f(x) \sim_c g(x)$  if  $f(x) \sim g(x)$  is conjectured but not proved. Likewise,  $=_c$  will denote a conjectured but unproved equality.

Oliveira e Silva [28, p. 2056] conjectured that

$$D_k(x) \sim_c k! x (\log x)^{k-1}, \qquad k \ge 1.$$
 (4)

The prime number theorem states:

$$\pi(x) \sim \frac{x}{\log x} \text{ as } x \to \infty \text{ or equivalently } p_n \sim n \log n \text{ as } n \to \infty.$$
 (5)

Because  $\pi(x) \sim x/\log x$  if and only if  $\log x \sim x/\pi(x)$ , and because  $x/\pi(x)$  is asymptotic to the average gap of the primes  $\leq x$ , the prime number theorem (5) implies that the average gap of the primes  $\leq x$  is asymptotic to  $\log x$ . (Why? Because *x* is asymptotic to the sum of gaps of the primes  $\leq x$ . Also  $\pi(x)$  is asymptotic to the number of gaps of the primes  $\leq x$ , by either definition of that number  $N_1$  or  $N_2$  above. By definition, the average of the gaps of the primes  $\leq x$  is the sum of gaps of the primes  $\leq x$  divided by the number of gaps of the primes  $\leq x$ . That ratio is asymptotic to  $x/\pi(x) = \log(x)$ .) As  $n \to \infty$ , (5) implies that  $p_n/n \to \infty$ . Taking logs of both sides gives  $\log p_n - \log n \to \infty$  but  $\log p_n/\log n \sim (\log n + \log \log n)/\log n \to 1$ .

Wolf [41] remarked that, when k = 0, the right side of conjecture (4) becomes  $x/\log x$  as in the prime number theorem in (5). When k = 1, (2) simplifies to  $D_1(x) = p_{\pi(x)} - 2$  and (4) asserts that  $D_1(x) \sim x$ , which is well known and, in any case, follows from Lemma 1. For k > 1, (4) apparently remains unproved, though well supported numerically [28, 41].

We shall use some elementary consequences of the prime number theorem (5).

262 👄 J. E. COHEN

Lemma 1. The prime number theorem (5) implies

$$\lim_{n \to \infty} \frac{p_{n+1}}{p_n} = 1,\tag{6}$$

$$\lim_{x \to \infty} \frac{x}{p_{\pi(x)}} = 1,\tag{7}$$

$$\lim_{x \to \infty} \frac{\log \pi(x)}{\log x} = 1.$$
(8)

**Proof.** By (5),

$$\frac{p_{n+1}}{p_n} \sim \frac{(n+1)\log(n+1)}{n\log n} \sim 1 \cdot 1 = 1.$$
(9)

This proves (6). To prove (7), observe that  $p_{\pi(x)} \le x < p_{\pi(x)+1}$ , hence  $1 \le x/p_{\pi(x)} < p_{\pi(x)+1}/p_{\pi(x)}$ . As  $x \to \infty$ , also  $\pi(x) \to \infty$ , so (6) implies (7). To prove (8), take logarithms of (5):  $\log \pi(x) \sim \log x - \log \log x \propto \log x$  as  $x \to \infty$ .

Now we define an exponentially distributed random variable and state, mostly without proof, several of its well known properties. Let *X* be a nonnegative real-valued random variable with cumulative distribution function (cdf)  $F(x) := \Pr\{X \le x\}, x \ge 0$ . If there exists some  $\lambda > 0$  such that  $F(x) = 1 - \exp(-\lambda x)$ , then *X* is *exponentially distributed* with scale parameter  $\lambda$  and we write  $X \stackrel{d}{=} Exp(\lambda)$ .

**Lemma 2.** Assume  $\lambda > 0$ . For n > 1, let X and  $X_1, \ldots, X_n$  be independently and identically distributed (iid)  $Exp(\lambda)$  random variables. *Then:* 

- 1. For any real c > 0,  $cX \stackrel{d}{=} Exp(\lambda/c)$ .
- 2. For real r > -1, the rth moment of X is  $E(X^r) = \Gamma(r+1)/\lambda^r$  [26, p. 28, Eq. (5)]. When  $r \in \mathbb{N}$ , then  $\Gamma(r+1) = r!$ . Hence the expectation or mean of X is  $1/\lambda$  and the variance of X is  $1/\lambda^2$ .
- 3. Let the order statistics of  $X_1, \ldots, X_n$  be  $X_{(1)} \leq \cdots \leq X_{(n)}$ . Then [30, p. 343] for  $i = 1, \ldots, n$ ,

$$E(X_{(i)}) = \frac{1}{\lambda} \sum_{j=1}^{i} \frac{1}{n-j+1} = \frac{1}{\lambda} \sum_{j=n-i+1}^{n} \frac{1}{j},$$
(10)

$$Var(X_{(i)}) = \frac{1}{\lambda^2} \sum_{j=1}^{i} \frac{1}{(n-j+1)^2} = \frac{1}{\lambda^2} \sum_{j=n-i+1}^{n} \frac{1}{j^2}.$$
(11)

4. The cumulative distribution function (cdf) of the smallest order statistic  $X_{(1)}$  is

$$\Pr\{X_{(1)} \le x\} = 1 - \exp(-n\lambda x), \ x \ge 0, \tag{12}$$

*i.e.*,  $X_{(1)} \stackrel{d}{=} Exp(\lambda n)$ . For 0 < q < 1, let  $y_1$  be the (1 - q)-quantile of  $X_{(1)}$ . By definition,  $y_1$  satisfies  $q = \Pr\{X_{(1)} > y_1\} = 1 - \Pr\{X_{(1)} \le y_1\} = \exp(-n\lambda y_1)$ . Then

$$y_1 = -(1/[n\lambda])\log q < +(1/[n\lambda])(1/q - 1).$$
(13)

In particular, if q = 1/n, then  $y_1 = +(1/[n\lambda]) \log n$ .

5. The cdf of the largest order statistic  $X_{(n)}$  is

$$\Pr\{X_{(n)} \le x\} = (1 - \exp(-\lambda x))^n, \ x \ge 0.$$
(14)

For 0 < q < 1, let  $y_n$  be the (1 - q)-quantile of  $X_{(n)}$ . By definition,  $y_n$  satisfies  $\Pr\{X_{(n)} > y_n\} = 1 - \Pr\{X_{(n)} \le y_n\} = q$ . Then  $y_n = -(1/\lambda)\log(1 - (1 - q)^{1/n}) < +(1/\lambda)(\log n - \log q).$  (15)

**Proof.**  $q = 1 - (1 - \exp(-\lambda y_n))^n$ , hence  $1 - \exp(-\lambda y_n) = (1 - q)^{1/n}$  and then  $-\lambda y_n = \log(1 - (1 - q)^{1/n})$ . The inequality follows from  $(1 - q)^{1/n} < 1 - q/n$ .

6. [18, p. 324, Ex. 9] By Lemma 2.1,  $\lambda X_n \stackrel{d}{=} Exp(1)$ . Then

$$\Pr\left\{\limsup_{n \to \infty} \frac{\lambda X_n}{\log n} = 1\right\} = 1.$$
(16)

7. Let  $S_n := X_1 + \dots + X_n$ . Let random variables  $U_1, \dots, U_{n-1}$  be iid uniformly on [0, 1] and let  $U_0 := 0$ ,  $U_n := 1$  with probability 1. Then  $(X_1/S_n, X_2/S_n, \dots, X_n/S_n) \stackrel{d}{=} (g_1, g_2, \dots, g_n) := (U_{(1)} - U_{(0)}, U_{(2)} - U_{(1)}, \dots, U_{(n)} - U_{(n-1)})$  ([15, p. 75, III.3(e)] and [18, p. 302, Ex. 42]).

8. [27] Large deviations: When  $a > 1/\lambda$ , then

$$\Pr\left\{\frac{X_1 + \dots + X_n}{n} > a\right\} \approx \exp(-(a\lambda - 1 - \log\lambda - \log a)n).$$
(17)

#### 3. Asymptotic moments of first *n* gaps are exponential

Define the *k*th moment (k > -1) of the first *n* gaps to be

$$\mu'_{k,n} := \frac{1}{n} \sum_{j=1}^{n} d_j^k.$$
(18)

Thus if  $\pi(x) = n$ , then (2) and (18) give  $\mu'_{k,n} = D_k(x)/n$ .

**Theorem 3.** The conjecture (4) of Oliveira e Silva [28, p. 2056] holds for each  $k \in \{0\} \cup \mathbb{N}$  if and only if the kth moment of the first n gaps is asymptotic to the kth moment of the exponential distribution  $Exp(1/\log n)$  with mean  $\log n$ :

$$\mu'_{k,n} \sim_c k! (\log n)^k, \qquad n \to \infty.$$
<sup>(19)</sup>

**Proof.** The cases k = 0, k = 1 are known to be true. Assume (4) for k > 1. Using the definition (2) and Lemma 1, replace x by  $p_{n+1}$  in (4) to get

$$D_k(p_{n+1}) \sim_c k! \, n \log n (\log(n \log n))^{k-1} = k! \, n \log n (\log n + \log \log n)^{k-1} \sim k! \, n (\log n)^k. \tag{20}$$

Dividing both sides by *n* gives (19).

Conversely, suppose the *k*th moment of the first *n* gaps is asymptotic to the *k*th moment of the exponential distribution  $Exp(1/\log n)$ . Then, working backward through the calculation in (20),

$$\mu_{k,n}' := \sum_{j=1}^{n} d_j^k \sim k! n (\log n)^k \sim k! n \log n (\log n + \log \log n)^{k-1} \\
 = k! n \log n (\log(n \log n))^{k-1} \sim_c D_k(p_{n+1}).$$
(21)

Again using Lemma 1 to replace  $p_{n+1}$  by x and  $n \log n \sim p_n \sim p_{n+1}$  by x gives

$$k! \, x (\log x)^{k-1} \sim_c D_k(x). \tag{22}$$

## 4. Asymptotic variance function obeys Taylor's power law

For  $n \in \mathbb{N}$ ,  $n \ge 2$ , define the mean and variance of the first *n* gaps as

$$m_n := \frac{1}{n} \sum_{j=1}^n d_j = \mu'_{1,n},\tag{23}$$

$$v_n := \frac{1}{n-1} \sum_{j=1}^n (d_j - m_n)^2 = \frac{n}{n-1} \left( \frac{1}{n} \sum_{j=1}^n d_j^2 - m_n^2 \right) = \frac{n}{n-1} (\mu'_{2,n} - (\mu'_{1,n})^2).$$
(24)

The central equality in (24) holds because  $\sum_{j=1}^{n} (d_j - m_n)^2 = \sum_{j=1}^{n} (d_j^2 - 2d_jm_n + m_n^2) = \sum_{j=1}^{n} d_j^2 - 2\sum_{j=1}^{n} d_jm_n + nm_n^2 = \sum_{j=1}^{n} d_j^2 - 2(nm_n)m_n + nm_n^2 = \sum_{j=1}^{n} d_j^2 - nm_n^2.$ 

The variance function of gaps is the function f that satisfies  $v_n = f(m_n)$  [1, 2, 11, 34–36]. The asymptotic variance function of gaps is a (not unique) function f that satisfies  $v_n \sim f(m_n)$ .

Corollary 4. Conditional on the conjectures of Heath-Brown [21], Wolf [39], and (4) of Oliveira e Silva [28, p. 2056],

$$\lim_{n \to \infty} \frac{\nu_n}{m_n^2} =_c 1.$$
<sup>(25)</sup>

**Proof.** From (19),

$$\frac{\mu'_{2,n}}{(\mu'_{1,n})^2} \sim_c 2.$$
(26)

Hence

$$v_n \sim (\mu'_{1,n})^2 \left(\frac{\mu'_{2,n}}{(\mu'_{1,n})^2} - 1\right) \sim_c (\mu'_{1,n})^2 (2-1) = m_n^2.$$
 (27)

This power-law asymptotic variance function is the special case c = 1, b = 2 of Taylor's law of fluctuation scaling  $v_n \sim cm_n^b$ , c > 0 [4–7, 12, 14, 33]. These parameter values c = 1, b = 2 are consistent with the exponential distribution, in which the variance equals the square of the mean. However, a variance asymptotic to the squared mean does not imply an exponential distribution (e.g., a normal distribution with mean  $\mu$  and variance  $\mu^2$  has a variance equal to the squared mean).

#### 5. An asymptotically exponential distribution of gaps is plausible

In a heuristic and numerical exploration, Yamasaki and Yamasaki [42] "assume that the exponential distribution can be applied to the gaps of prime numbers. But the gaps [except for  $d_1 = 1$ ] are always even integers, so do not distribute continuously on  $[0, \infty)$ ." They proposed adjustments to the exponential distribution.

Conditional on a uniform version of the unproved prime *r*-tuple conjecture of Hardy and Littlewood [20], Gallagher [16] showed that, for  $\lambda > 0$ ,  $h \sim \lambda \log N$ , and  $n \leq N$ , as  $N \to \infty$ , the distribution of the values of  $\pi(n + h) - \pi(n)$  converges to the Poisson distribution with parameter  $\lambda$ . Thus  $\lambda$  is the expectation and the variance of the number of primes over the interval  $(n, n + \lambda \log N]$  of length  $\lambda \log N$ , so the average gap is asymptotically  $\log N$ . If the Poisson-distributed number of primes arrived in the interval  $(n, n + \lambda \log N]$  according to a Poisson process, then the intervariation interval between successive primes (that is, the gap) would be exponentially distributed [15, pp. 188, 378],  $Exp(1/\log N)$ .

Wolf [40] gave strong numerical evidence [40, p. 3, his Figure 1] and a heuristic argument [40, p. 2, his Eq. (7)] that an exponential distribution with mean log *n* asymptotically approximates the distribution of the first *n* gaps. To circumvent the difficulty that the exponential distribution lacks the discreteness of the natural numbers, Wolf [40, his Eq. (19), his Figure 5] proposed a rescaling that turns the discrete  $d_n$  into a continuous variable.

Our finding that, conditional on the conjecture (4) of Oliveira e Silva [28, p. 2056], the moments of the first *n* gaps are asymptotic to the moments of the exponential distribution, even though the gaps are not continuously distributed, is consistent with T. Stieltjes's discovery in 1894 that the moments of the lognormal distribution do not uniquely specify the continuous probability distribution that produced the moments [31, pp. 17–18, section 2.1].

We suggest it is plausible that the moments (18) of the first *n* gaps are asymptotic to the moments of  $Exp(1/\log n)$ . First, the primes that do not exceed *x* are asymptotically uniformly distributed on [0, x] as  $x \to \infty$  [4]. To see why, choose any 0 < r < 1. Then, for large *x*, the number of primes not exceeding *rx* divided by the number of primes not exceeding *x* is asymptotically, by the prime number theorem (5),  $\pi(rx)/\pi(x) \sim \lim_{x\to\infty} (rx/\log(rx))/(x/\log x) = r$ . (From a more general perspective, the asymptotic counting function  $\pi(x) \sim x/\log x$  of primes is regularly varying [3] with exponent 1.)

Assume  $U_1, \ldots, U_n$  are iid uniform on [0, x]. The finding that the primes are asymptotically uniformly distributed on [0, x] [4] suggests the order statistics  $U_{(1)} \leq \cdots \leq U_{(n)}$  of  $U_1, \ldots, U_n$  as a stochastic model of the asymptotic distribution of the primes on [0, x] and suggests the *n* "spacings" [5, 22]  $g_i := U_{(i)} - U_{(i-1)}$ ,  $i = 1, \ldots, n$ , with  $U_{(0)} := 0$  almost surely, as a stochastic model of the first *n* gaps. The spacings are exponentially distributed (Lemma 2.7) and their average size is asymptotically log *n*. So the gaps between consecutive primes  $\leq x$  are plausibly distributed as the spacings between consecutive order statistics of  $\pi(x)$  iid uniform random variables on [0, x], and these spacings are distributed as  $Exp(1/\log n)$ .

This stochastic model of the primes and gaps intentionally omits the discreteness of integers and the lack of independence of primes.

## 6. The largest of the first n gaps

What would the exponential model of gaps predict about the asymptotic behavior of the maximal gap,  $G_n := \max\{d_1, \ldots, d_n\}$ , among the first *n* gaps? Here we consider two possible answers.

**Theorem 5.** For fixed n > 1, let  $X_i \stackrel{d}{=} Exp(1/\log n)$ , i = 1, ..., n, be iid. Let  $\gamma \approx 0.5772$  be the Euler-Mascheroni constant,  $\zeta(\cdot)$  the zeta function, and  $\pi^2/6 \approx 1.6449$ . Then the mean and variance of the largest order statistic  $X_{(n)}$  are

$$E(X_{(n)}) = (\log n) \sum_{j=1}^{n} \frac{1}{n-j+1} \sim (\log n)(\gamma + \log n) \sim (\log n)^{2},$$
(28)

$$Var(X_{(n)}) = (\log n)^2 \sum_{j=1}^n \frac{1}{(n-j+1)^2} \sim (\log n)^2 \zeta(2) = (\log n)^2 \frac{\pi^2}{6}.$$
(29)

Thus  $Var(X_{(n)})/E(X_{(n)}) \rightarrow \pi^2/6 > 1$  and  $Var(X_{(n)}) \sim (\pi^2/6)E(X_{(n)})$ . The maximal gap illustrates Taylor's law with  $c = \pi^2/6$ , b = 1.

**Proof.** From Lemma 2.3,

$$E(X_{(n)}) = (\log n) \left( 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{n} \right) \sim (\log n)(\gamma + \log n) \sim (\log n)^2,$$
(30)

$$Var(X_{(n)}) = (\log n)^2 \left( 1 + \frac{1}{2^2} + \frac{1}{3^2} + \dots + \frac{1}{n^2} \right) \sim (\log n)^2 \zeta(2) = (\log n)^2 \frac{\pi^2}{6}.$$
(31)

The Cramér-Shanks conjecture may be stated as

$$\max_{p_j \le x} (p_{j+1} - p_j) \sim_c (\log x)^2 \sim (\log p_{\pi(x)})^2.$$
(32)

See Cramér [8, pp. 24, 27], Shanks [32, p. 648, his Eq. (5)], and the very helpful Granville [17, p. 21, his Eq. (14)].

**Theorem 6.** The Cramér-Shanks conjecture (32) holds if and only if the maximal gap  $G_n := \max\{d_1, \ldots, d_n\}$  satisfies

$$G_n \sim_c (\log n)^2. \tag{33}$$

**Proof.** If there are *n* primes less than or equal to *x*, then the largest of these is  $p_n \sim n \log n \sim x$ , using Lemma 1. Thus, translated from the scale of primes not exceeding *x* to the first *n* gaps, the Cramér-Shanks conjecture (32) becomes (33) because

$$\max_{1 \le j \le n} d_j \sim_c (\log(n \log n))^2 = (\log n + \log \log n)^2 \sim (\log n)^2.$$
(34)

Conversely, if  $\max_{1 \le j \le n} d_j \sim_c (\log n)^2$ , then  $\max_{1 \le j \le n} d_j = \max_{p_j \le p_n} d_j \sim \max_{p_j \le x} d_j$  and  $\log x \sim \log(n \log n) \sim \log n + \log \log n$ , so  $(\log x)^2 \sim (\log n)^2$  and (32) holds.

Theorems 5 and 6 suggest that  $(\log x)^2$  in the Cramér-Shanks conjecture (32) is too small asymptotically. In the model of Theorem 5, the largest gap is likely to be larger than the expectation of the largest gap because the variance of the largest gap is of the same order of magnitude as the expectation of the largest gap. To the extent that the above conjectures are correct that gaps have exponential moments asymptotically, it would be surprising if the largest of the first *n* gaps did not exceed  $(\log x)^2$ .

These suggestions indirectly challenge Firoozbakht's conjecture [38]. Kourbatov [23] showed that Firoozbakht's conjecture implies that  $d_n := p_{n+1} - p_n <_c (\log p_n)^2 - \log p_n - 1$  for all n > 9 and is implied by  $d_n <_c (\log p_n)^2 - \log p_n - 1.17$  for all n > 9. These upper bounds are stronger than the Cramér-Shanks conjecture (32), and lower than the dominant term  $(\log p_n)^2 \sim (\log n)^2$  (see Theorem 6) by amounts that increase without limit as *n* increases. To the extent that our Theorems 5 and 6 suggest that  $(\log x)^2$  in the Cramér-Shanks conjecture (32) is too small asymptotically, our results suggest even more that Kourbatov's upper bounds on gaps are likely to be too small. Kourbatov [23] discusses many related inequalities and conjectures concerning prime gaps.

Granville [17, p. 24] reviewed results that suggest, contrary to the Cramér-Shanks conjecture (32), that

$$\max_{p_j \le x} d_j \stackrel{>}{\sim}_c 2e^{-\gamma} (\log x)^2 \sim 2e^{-\gamma} (\log p_{\pi(x)})^2,$$
(35)

where  $2e^{-\gamma} \approx 1.1229$ . Translating from the scale of *x* to the scale of the number of gaps  $n = \pi(x)$  by  $x \sim p_n \sim n \log n$  so that  $(\log x)^2 \sim (\log(n \log n))^2 \sim (\log n)^2$  gives the proposal that

$$G_n \stackrel{>}{\sim}_c 2e^{-\gamma} (\log n)^2. \tag{36}$$

These observations motivate another estimate of the largest of the first *n* gaps, namely, the (1 - 1/n)-quantile of the largest order statistic  $X_{(n)}$ . We now determine the value of *y* such that, in a sample of size *n* from  $Exp(1/\log n)$ , as in Theorem 5, the probability that the largest order statistic  $X_{(n)}$  exceeds *y* is 1/n.

**Theorem 7.** In a sample of size k > 1 from  $Exp(1/\log n)$ , let  $y_k$  be the (1 - 1/k)-quantile of the maximal order statistic  $X_{(k)}$ . Then  $y_k \sim 2(\log n)(\log k)$  as  $k \to \infty$ . If  $k \sim tn$  for some t > 0, then  $y_k \sim 2(\log n)^2$  as  $n \to \infty$ .

**Proof.** Using Lemma 2.5, (15), with  $\lambda = 1/\log n$  gives

$$y_k = -(\log n)\log(1 - (1 - 1/k)^{1/k}).$$
(37)

Repeated applications of l'Hopital's rule as  $k \to \infty$  shows that  $-\log(1 - (1 - 1/k)^{1/k}) \sim (2\log k)$  and hence that

$$y_k \sim +2(\log n)(\log k). \tag{38}$$

Wolf [40, his Eq. (15)] argued heuristically for the conjecture that, as  $x \to \infty$ ,

$$G_W(x) := \max_{p_n < x} (p_n - p_{n-1}) \sim_c \frac{x}{\pi(x)} (2\log(\pi(x)) - \log(x) + c), \tag{39}$$

where  $c \approx 0.2778769$ . Here  $c := \log(C_2)$  and  $C_2 := 2 \prod_{p>2} (1 - (p-1)^{-2})$  is known sometimes as the twins constant and sometimes as twice the twins constant. The product in  $C_2$  is taken over all positive odd primes. Replacing *x* by  $p_n + \varepsilon$  and letting  $\varepsilon \downarrow 0$  gives the equivalent conjecture, as  $n \to \infty$ ,

$$G_W(p_n) \sim_c \frac{p_n}{n} (2\log(n) - \log(n\log n) + c) \sim (\log n)^2.$$
 (40)

# 7. Numerical illustrations

For moments and maximal gaps, we compare exact numerical results derived from public sources with asymptotic expressions derived from the exponential model of prime gaps and from other conjectures.

#### 7.1. Moments

To illustrate Theorem 3, we compare exact computations of the moments of the first *n* gaps with the moments of an exponential distribution with mean log *n* as given by (19). Wolf [40] generously made public extensive numerical results at http://pracownicy. uksw.edu.pl/mwolf/gapstau.zip. Wolf [40, p. 2, his Eq. (2)] defined, for large real *x* and *even*  $d \in \mathbb{N}$ ,

$$\tau_d(x) := \#\{p_{n+1} \mid p_{n+1} < x \text{ and } p_{n+1} - p_n = d\}$$
(41)

and, for every *odd*  $d \in \mathbb{N}$ ,  $\tau_d(x) := 0$ . In words, when *d* is even,  $\tau_d(x)$  is the number of pairs of consecutive primes such that the larger prime is less than *x* and such that their difference equals *d*; and for odd *d*,  $\tau_d(x) := 0$ . This definition and my subsequent analyses exclude  $\tau_1(x) = 1$ ,  $x \ge 3$ , representing the single gap  $p_2 - p_1 = 3 - 2$  of size d = 1. Wolf collected the non-zero values of  $\tau_d(x)$  in 34 text files, one file for each of the 34 values of  $x = 2^t$ , t = 15(1)48. (Approximately,  $2^{48} \approx 2.8147 \times 10^{14}$ .) Each file has two columns, the first listing values of *d* such that  $\tau_d(x) > 0$ , and the second listing the corresponding non-zero values of  $\tau_d(x)$ .

First, using MATLAB Version 9.13.0.2049777 (R2022b), I calculated  $\tau_d(x)$  directly for all primes less than  $2^{20}$ . My results agreed exactly with those in Wolf's file tau20s.dat. For example, Wolf and I independently found  $\tau_2(2^{20}) = 8535$  gaps of size d = 2; the maximal gap size, d = 114, occurred once:  $\tau_{114}(2^{20}) = 1$ .

Having partially verified Wolf's results, I analyzed Wolf's 12 files with values of  $x = 2^t$ , t = 15, 18, 21, 24, 27, ..., 48. For each such file separately, Table 1 gives *t*; the number of gaps  $n = \sum_{d \in \mathbb{N}} \tau_d(x)$  where  $x = 2^t$  (not counting the first gap d = 1); the first four integer moments  $\mu'_{k,n}$ , k = 1, 2, 3, 4 from (18); and the maximal gap  $G_n$ . Figure 1 compares  $\mu'_{k,n}$  with  $k!(\log n)^k$ , which is asymptotic

**Table 1.** For each upper limit  $x = 2^t$ , this table shows the exponent t of 2, the number n of gaps between consecutive primes (not counting the odd first gap), the moments  $\mu'_{k,n}(k = 1, 2, 3, 4)$ , and the maximal gap  $G_n$ .

			Moments				
row	t	n	k = 1	<i>k</i> = 2	<i>k</i> = 3	k = 4	Gn
1	15	3510	9.3293	136.2017	2.7818e+03	7.4292e+04	72
2	18	22998	11.3982	210.7095	5.5060e+03	1.8546e+05	86
3	21	155609	13.4770	304.1124	9.8914e+03	4.2503e+05	148
4	24	1077869	15.5652	412.7866	1.5776e+04	7.8862e+05	154
5	27	7603551	17.6520	539.4491	2.3885e+04	1.3864e+06	222
6	30	54400026	19.7379	683.2373	3.4423e+04	2.2806e+06	282
7	33	393615804	21.8231	844.1273	4.7670e+04	3.5440e+06	354
8	36	2.8744e+09	23.9074	1.0222e+03	6.3972e+04	5.2773e+06	464
9	39	2.1152e+10	25.9908	1.2173e+03	8.3638e+04	7.5819e+06	532
10	42	1.5666e+11	28.0736	1.4296e+03	1.0699e+05	1.0574e+07	652
11	45	1.1667e+12	30.1560	1.6590e+03	1.3435e+05	1.4377e+07	766
12	48	8.7312e+12	32.2379	1.9056e+03	1.6603e+05	1.9127e+07	906



**Figure 1.** Moments  $\mu'_{k,n}$  (k = 1, 2, 3, 4) of the first *n* gaps between consecutive primes (solid dots) from k = 1 (bottom row) to k = 4 (top row); and corresponding *k*th moments (k = 1, 2, 3, 4) of exponential distributions *Exp*(1/log *n*) (solid lines). The dots and the lines are calculated independently with no adjustment of parameters.



**Figure 2.** Maximal gap  $G_n$  (solid blue dots) among the first n gaps and four conjectured models:  $(\log n)^2$  (solid black line);  $(\log p_n)^2$  (solid black

**Table 2.** Seven maximal gaps  $G_n$  that exceed  $(\log n)^2$  and  $2e^{-\gamma} (\log n)^2$ , the index *n* of the prime  $p_n$  that begins the maximal gap, the prime  $p_n$  that begins the maximal gap,  $(\log p_n)^2$ , and  $2e^{-\gamma} (\log p_n)^2$ .

row	п	Gn	p <sub>n</sub>	$(\log n)^2$	$(\log p_n)^2$	$2e^{-\gamma}(\log n)^2$	$2e^{-\gamma}(\log p_n)^2$
1	1	1	2	0	0.4805	0	0.5395
2	2	2	3	0.4805	1.2069	0.5395	1.3553
3	4	4	7	1.9218	3.7866	2.1580	4.2520
4	9	6	23	4.8278	9.8313	5.4212	11.0398
5	30	14	113	11.5681	22.3482	12.9901	25.0952
6	217	34	1327	28.9433	51.7058	32.5010	58.0614
7	49749629143526	1132	1693182318746371	994.6470	1229.6	1116.9	1380.7

Only for n = 1 and n = 2 is  $G_n > 2e^{-\gamma} (\log p_n)^2$ .

to the corresponding moments of samples of size *n* from an exponential distribution  $Exp(1/\log n)$ , as conjectured in (19). Visually, the agreement in Figure 1 between the exact moments and the asymptotic moments of  $Exp(1/\log n)$  is remarkable.

## 7.2. Maximal gaps

We compare the first 80 known maximal gaps (now *including* the initial gap of size 1) [37] with four asymptotic models: (33), (32), (36), and (35). These results on maximal gaps are more extensive and more detailed than those in Table 1. The tabulation lists three sequences:

- 1. the indices of primes after which a maximal gap occurs
- $n = 1, 2, 4, 9, 24, 30, 99, 154, 189, 217, 1183, \dots$  (OEIS <u>A005669</u>),
- 2. the corresponding maximal gaps
- $G_n := p_{n+1} p_n = 1, 2, 4, 6, 8, 14, 18, 20, 22, 34, 36, \dots$  (OEIS <u>A005250</u>), and
- 3. the primes after which a maximal gap occurs
  - $p_n = 2, 3, 7, 23, 89, 113, 523, 887, 1129, 1327, 9551, \dots$  (OEIS <u>A002386</u>).

I partially verified the Wikipedia tabulation in two ways. First, Marek Wolf (personal communication, 2022-11-10) generously sent me his tabulation of the first 74 maximal gaps  $G_n$  (not including  $G_1$ ), their indices n, and their beginning primes  $p_n$ . Second, using MATLAB, I calculated these three sequences directly for all primes less than 10<sup>6</sup>. My results and Wolf's agreed exactly, as far as they went, with the Wikipedia tabulation [37].

Figure 2 plots, as a function of *n*, the maximal gap size  $G_n$  among the first *n* gaps; and the values of the four conjectured asymptotic expressions  $(\log n)^2$ ;  $(\log p_n)^2$ ;  $2e^{-\gamma} (\log n)^2$ , where  $2e^{-\gamma} \approx 1.1229$ ; and  $2e^{-\gamma} (\log p_n)^2$ . The majority of points  $(n, G_n)$  (blue solid dots in Figure 2) fall slightly below the conjectured asymptotic behavior  $(n, (\log n)^2)$  (solid black line in Figure 2) derived here from the conjectured asymptotically exponential distribution of gaps. The form of Wolf's conjecture in (40) falls slightly below  $(n, (\log n)^2)$  and often closer to the points  $(n, G_n)$  than  $(n, (\log n)^2)$ . However, the seven exceptional values of *n* with  $G_n > 2e^{-\gamma} (\log n)^2$  in Table 2 suggest that  $\limsup_{n\to\infty} G_n/[2e^{-\gamma} (\log n)^2]$  may exceed 1. The available numerical results neither confirm nor reject the

suggestion implied by Theorem 7 that  $\limsup_{n\to\infty} G_n$  might be asymptotic to  $2(\log n)^2$ . The asymptotic behavior of  $G_n$  remains to be determined.

# Acknowledgments

J.E.C. thanks Andrew Granville, D. R. Heath-Brown, Tomás Oliveira e Silva, and Marek Wolf for very helpful guidance to relevant references; Marek Wolf for sharing numerical results analyzed here; and Roseanne Benjamin for assistance during this work. Two helpful reviewers provided constructive suggestions.

# **Declaration of Interest**

The author reports there are no competing interests to declare.

# ORCID

Joel E. Cohen D http://orcid.org/0000-0002-9746-6725

# References

- [1] Bar-Lev, S. K., Enis, P. (1986). Reproducibility and natural exponential families with power variance functions. Ann. Stat. 14(4): 1507–1522.
- [2] Bar-Lev, S. K., Stramer, O. (1987). Characterizations of natural exponential families with power variance functions by zero regression properties. *Probab. Theory Related Fields* 76: 509–522.
- [3] Bingham, N. H., Goldie, C. M., Teugels, J. L. (1987). Regular Variation, Encyclopedia of Mathematics and its Applications. Cambridge, England: Cambridge University Press.
- [4] Cohen, J. E. (2016). Statistics of primes (and probably twin primes) satisfy Taylor's law from ecology. Amer. Stat. 70(4): 399-404.
- [5] Cohen, J. E. (2020). Species-abundance distributions and Taylor's power law of fluctuation scaling. Theor. Ecol. 13(4): 607-614.
- [6] Cohen, J. E. (2023). Integer sequences with regularly varying counting functions have power-law variance functions. Integers 23(A87): 1–19.
- [7] Cohen, M. P. (2017). Non-asymptotic mean and variance also approximately satisfy Taylor's law. Amer. Stat. 71(2): 187.
- [8] Cramér, H. (1936). On the order of magnitude of the difference between consecutive prime numbers. Acta Arith. 2(1): 23-46.
- [9] Crandall, R., Pomerance, C. (2005). Prime Numbers-A Computational Approach, 2nd ed. New York: Springer.
- [10] Cutter, P. A. (2001). Finding prime pairs with particular gaps. *Math. Comp.* 70(236): 1737–1744.
- [11] Davidian, M., Carroll, R. J. (1987). Variance function estimation. J. Amer. Stat. Assoc., 82(400): 1079–1091.
- [12] Demers, S. (2018). Taylor's law holds for finite OEIS integer sequences and binomial coefficients. Amer. Stat. 72(4): 376–378.
- [13] Dickson, L. G. (1919). History of the Theory of Numbers, Volume I: Divisibility and Primality, Vol. 256(1). Washington, DC: Carnegie Institution of Washington.
- [14] Eisler, Z., Bartos, I., Kertész, J. (2008). Fluctuation scaling in complex systems: Taylor's law and beyond. Adv. Phys. 57(1): 89-142.
- [15] Feller, W. (1971). An Introduction to Probability Theory and Its Applications, Vol. 2, 2nd ed. New York: Wiley.
- [16] Gallagher, P. X. (1981). On the distribution of primes in short intervals. Mathematika 23(1): 4-9. Corrigendum, Mathematika 28: 86.
- [17] Granville, A. (1995). Harald Cramér and the distribution of prime numbers. Scand. Actuar. J. 1: 12-28.
- [18] Grimmett, G. R., Stirzaker, D. R. (2001). Probability and Random Processes, 3rd ed. New York: Oxford University Press.
- [19] Guy, R. K. (1994). Unsolved Problems in Number Theory, 2nd ed. New York: Springer Verlag.
- [20] Hardy, G. H., Littlewood, J. E. (1966). Some problems of 'partitio numerorum' : III: On the expression of a number as a sum of primes. Acta Math. 44:1-70. Reprinted in "Collected Papers of G. H. Hardy," Vol. I, pp. 561–630. Oxford: Clarendon Press.
- [21] Heath-Brown, D. R. (1992). Gaps between primes and the pair correlation of zeros of the zeta-function. Acta Arith. 41: 85-99.
- [22] Holst, L. (1980). On the lengths of the pieces of a stick broken at random. J. Appl. Probab. 17: 623-634.
- [23] Kourbatov, A. (2015). Upper bounds for prime gaps related to Firoozbakht's conjecture. arXiv 1506.03042v4.
- [24] Maier, H. (1981). Chains of large gaps between consecutive primes. Adv. Math. 39(3): 257-269.
- [25] Maier, H. (1985). Primes in short intervals. Michigan Math. J. 32(2): 221-225.
- [26] Marshall, A. W., Olkin, I. (2007). Life Distributions: Structure of Nonparametric, Semiparametric, and Parametric Families. Series in Statistics. New York: Springer.
- [27] Massachusetts Institute of Technology. (2013). Large deviations for i.i.d. random variables, Advanced Stochastic Processes. Available at: https://ocw.mit. edu/courses/15-070j-advanced-stochastic-processes-fall-2013/ed35842f2e902421e4a6cdd4d9eca6fe\_MIT15\_070JF13\_Lec2.pdf. [Online; accessed 13-April-2024].
- [28] Oliveira e Silva, T., Herzog, S., Pardi, S. (2014). Empirical verification of the even Goldbach conjecture and computation of prime gaps up to  $4 \times 10^{18}$ . *Math. Comp.* 83(288): 2033–2060.
- [29] Riesel, H. (1994). Prime Numbers and Computer Methods for Factorization. Progress in Mathematics, Vol. 126, 2nd ed. Boston: Birkhauser.
- [30] Sarhan, A. E., Greenberg, B. G., eds. (1962). Contributions to Order Statistics. New York, London: Wiley.
- [31] Schmüdgen, K. (2020). Ten lectures on the moment problem. Available at: https://doi.org/10.48550/arXiv.2008.12698. arXiv:2008.12698v1 [math.FA].
- [32] Shanks, D. (1964). On maximal gaps between successive primes. Math. Comp. 18(88): 646-651.
- [33] Taylor, R. A. J. (2019). Taylor's Power Law: Order and Pattern in Nature. New York: Academic Press.
- [34] Tweedie, M. C. K. (1946). The regression of the sample variance on the sample mean. J. London. Math. Soc. 21: 22-28.
- [35] Tweedie, M. C. K. (1947). Functions of a statistical variate with given means, with special reference to Laplacian distributions. *Proc. Cambridge Philos.* Soc. 43: 41–49.
- [36] Tweedie, M. C. K. (1984). An index which distinguishes between some important exponential families. In: Ghosh, J. K., Roy, J., eds. Statistics: Applications and New Directions. Proceedings of the Indian Statistical Institute Golden Jubilee International Conference, pp. 579–604, Calcutta, 1984. Indian Statistical Institute.
- [37] Wikipedia contributors. (2022). Prime gap Wikipedia, the free encyclopedia. Available at: https://en.wikipedia.org/w/index.php?title= Prime\_gap&oldid=1113313971. [Online; accessed 13-November-2022].
- [38] Wikipedia contributors. (2023). Firoozbakht's conjecture Wikipedia, the free encyclopedia. https://en.wikipedia.org/w/index.php?title=Firoozbakht %27s\_conjecture&oldid=1167645599. [Online; accessed 10-April-2024].

- [39] Wolf, M. (1998). Some conjectures on the gaps between consecutive primes. Available at: https://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1. 49.3015&rep=rep1&type=pdf and http://pracownicy.uksw.edu.pl/mwolf/conjectures.ps, 1998. Preprint accessed 2022-09-23.
- [40] Wolf, M. (2014). Nearest-neighbor-spacing distribution of prime numbers and quantum chaos. *Phys. Rev. E* 89: 022922.
- [41] Wolf, M. (2017). On the moments of the gaps between consecutive primes. arXiv:1705.10766v1.
- [42] Yamasaki, Y., Yamasaki, A. (1994). On the gap distribution of prime numbers. Notes Res Inst. Math. Anal. Kyoto Univ. 887: 151-168.