

## Origin of patterning in neural tubes

SIR — A notochord and dorsal tubular nerve cord are essential features of chordates, and must have evolved in the most ancient members of our phylum. How complex was the primitive chordate neural tube? Ascidians, as members of the urochordate subphylum, derived from the most basal extant lineage within Chordata<sup>1</sup>; anatomically, the nerve cord of ascidian tadpole larvae is simpler than that of vertebrates or cephalochordates, with less than 100 rostral neurons and just axons and ependymal cells in the tail<sup>2</sup>. It is unclear whether this reflects a simple ancestral pattern, or whether it masks hidden developmental complexity.

Here we report the expression of an ascidian (*Halocynthia roretzi*) *Pax* gene,

*HrPax-37*, descendant from the precursor of *Pax-3* and *Pax-7* in vertebrates. The last two are expressed dorsally in both neural tube and somitic mesoderm; in addition, mouse *Pax-3* mutations reveal a function in differentiation of the dorsal neural tube<sup>3-5</sup>. Expression of *HrPax-37* is first detected at the early gastrula stage (*a* in the figure) in the nuclei of six bilateral pairs of cells, destined to form the dorsal part of the neural tube (a8.25, b8.19 and b8.17; the last two also yield muscle cells) or dorsal epidermis (a8.26, b8.20 and b8.18)<sup>6,7</sup>. By the mid-gastrula stage, expression is maintained only in the first three cells (*b* in the figure). In contrast, at the later neural-plate stage the expression becomes downregulated in the primordial

dorsal nerve cells and reactivated in the dorsal epidermis (*c* in the figure). At the neurula stage, just a single row of epidermal cells flanking the presumptive neural tube continues to express the gene (*d* in the figure).

Expression during these early embryonic stages indicates that *HrPax-37* might have a similar function to vertebrate *Pax-3* and *Pax-7* in the differentiation of dorsal neural tube, although *HrPax-37* is also expressed in dorsal epidermis. Epidermal sensory neurons of ascidian larvae are derived from the dorsal epidermal cells<sup>7,8</sup>, raising the possibility that the expression in this region is comparable with *Pax-3* in vertebrate neural crest<sup>3</sup>.

During closure of the neural tube, expression fades from dorsal epidermal cells, but is reactivated in the three populations of cells in the neural tube (*e, f* in the figure). By the tadpole larva stage, expression in the most rostral cell population disappears, although extensive expression persists in the sensory vesicle and visceral ganglion (*g* in the figure). At this stage, a striking pattern of *HrPax-37* expression emerges more posteriorly, in the nerve cord of the tail. Expression is detected in roughly 15 reiterated spots, within a dorsal row of cells in the nerve cord (*h* in the figure). To our knowledge,

no segmental structures have previously been reported in the ascidian neural tube. Because this expression must derive from ependymal cells (there are no neuronal cell bodies in the nerve cord<sup>2</sup>), we suggest this reiterated distribution of *HrPax-37* is an evolutionary remnant reflecting ancestry from a more elaborate segmental organization. This suggestion is compatible with the existence of a segmental ancestor for all urochordates, and would resolve the paradox that appendicularians (also urochordates) have motor neurons segmentally distributed in the tail neural tube<sup>9,10</sup>.

In conclusion, we suggest that dorsal specification by genes of the *Pax-3/7* subfamily, and segmental organization of neural tube, were established in the ancestors of extant chordates during emergence of the dorsal tubular nervous system.

**Hiroshi Wada**

**Peter W. H. Holland**

School of Animal and Microbial Sciences,  
University of Reading, Whiteknights,  
Reading RG6 6AJ, UK

**Noriyuki Satoh**


Department of Zoology,  
Graduate School of Sciences,  
Kyoto University, Kyoto 606-01, Japan

1. Wada, H. & Satoh, N. *Proc. Natl. Acad. Sci. USA* **91**, 1801–1804 (1994).
2. Nicol, D. & Meinaertzhagen, I. A. *J. Comp. Neurol.* **309**, 415–429 (1991).
3. Goulding, M. D., Chalepakis, G., Deutsch, U., Erselius, J. R. & Gruss, P. *EMBO J.* **10**, 1135–1147 (1991).
4. Jostes, B., Walther, C. & Gruss, P. *Mech. Dev.* **33**, 27–38 (1991).
5. St-Onge, L., Pituello, F. & Gruss, P. *Semin. Dev. Biol.* **6**, 285–292 (1995).
6. Conklin, E. G. *J. Acad. Nat. Sci. (Philadelphia)* **13**, 1–119 (1905).
7. Nishida, H. *Dev. Biol.* **121**, 526–541 (1987).
8. Torrence, S. A. & Cloney, R. A. *Zoomorphology* **99**, 103–115 (1982).
9. Flood, P. R. *J. Microsc.* **18**, 317–326 (1973).
10. Bone, Q. *Am. Zool.* **29**, 5–18 (1989).

## Neuronal cell death and tPA

SIR — The serine protease tissue plasminogen activator (tPA) has been implicated in brain hippocampal function, as its messenger RNA is present<sup>1</sup> and is rapidly induced in the rat and mouse hippocampus on pharmacological or electrical induction of neuronal activity<sup>2,3</sup>. In addition, mice deficient in tPA (tPA<sup>-/-</sup>)<sup>4</sup> are resistant to neuronal destruction<sup>5</sup> after intra-hippocampal injection of high concentrations of glutamate analogues (excitotoxins<sup>6,7</sup>). Here we investigate whether tPA<sup>-/-</sup> mice are resistant to neuronal degeneration because of pre- and/or postnatal development in the absence of tPA, or whether tPA is necessary for neuronal death in the adult mouse.

An adult requirement for tPA would be significant for two reasons. First, tPA has recently been approved for treatment



Spatial and temporal expression of *HrPax-37* as deduced from *in situ* hybridization. *a*, Lateral view of embryo at early gastrula stage. The signal of b8.17 is out of the plate of focus. *b*, Dorsal view of mid-gastrula embryo. Anterior is to the top in *a* and *b*. *c, d*, Dorsal view of embryo at neural-plate stage (*c*) and neurula stage (*d*). *e*, Dorsal view of the embryo at initiating tailbud stage. The neural tube is closing from the posterior region; in more anterior regions, the expression of *HrPax-37* remains in epidermis. *f*, Lateral view of the embryo at mid-tailbud stage. Anterior is to the right in *c-f*. *g*, Dorsal view of the trunk region of larva. s: sensory vesicle, v: visceral ganglion. *h*, Lateral view of larval tail. The segmental expression of *HrPax-37* in the nerve cord is indicated by arrowheads. Arrows, dorsal and ventral boundary of the neural tube; n, notochord; nt, neural tube. Anterior is to the left in *g* and *h*.

of thrombotic stroke. Because the blood-brain barrier is compromised during stroke, tPA might gain access to the brain parenchyma, thus contributing to neuronal cell death. Second, inhibition of tPA activity in the adult mouse could protect against certain neuronal pathologies.

To distinguish between a developmental or an acute need for tPA, we delivered tPA protein bilaterally to the hippocampus of tPA<sup>-/-</sup> mice, then injected kainate (a glutamate analogue) on one side, and assessed the extent of neuronal

death. We found that tPA<sup>-/-</sup> mice infused with buffer (control) did not exhibit significant neuronal degeneration after kainate injection (Fig. 1, top)<sup>5</sup>. In contrast, tPA<sup>-/-</sup> mice infused with tPA exhibited dramatic neurodegeneration in the kainate-injected hippocampus (Fig. 1, bottom). We detected proteolytic activity<sup>1</sup> equally over both sides of the hippocampus of the tPA-infused tPA<sup>-/-</sup> mice (verfying that tPA was delivered appropriately), whereas we observed no tPA activity in the control tPA<sup>-/-</sup> mice. Because we observed degeneration only in the kainate-injected side, tPA alone at this concentration is not sufficient to kill neurons. These results indicate that tPA is required for neuronal death in the adult animal at the time of excitotoxic insult.

As tPA is required to promote neuronal degeneration, we tested whether inhibition of its enzymatic activity might retard neuronal death. We infused mice with plasminogen activator inhibitor-1 (PAI-1; a serine-protease inhibitor that inhibits tPA<sup>8</sup>), injected them with kainate, and assessed neuronal survival in the hippocampus. Wild-type mice infused with buffer were sensitive to neuronal degeneration in the hippocampus (Fig. 2, top). In contrast, mice infused with PAI-1 were resistant to kainate-induced neuronal death (Fig. 2, bottom). The resistance of wild-type mice infused with PAI-1 was comparable to that observed with buffer-infused tPA<sup>-/-</sup> mice.

Various neuropathological conditions may involve excitotoxic damage to the brain<sup>9</sup>. The identification of extracellular proteases as contributing to the degeneration pathway provides an attractive target for therapeutic intervention. One concern of such therapy in humans would be that inhibition of tPA might lead to clotting disorders. However, tPA<sup>-/-</sup> mice do not exhibit spontaneous thromboses<sup>4</sup>, indicating that blood homeostasis is possible in the absence of tPA. Therefore, it seems reasonable to investigate further the

effectiveness of protease inhibitors for the therapy of excitotoxic-mediated brain disorders.

Thrombotic stroke is thought to involve an excitotoxic pathway<sup>9</sup>, and tPA can promote neuronal death after excitotoxic insult. Recent studies report treatment of ischaemic stroke within three hours of onset with intravenous tPA<sup>10,11</sup>. Certainly the excitotoxic cell death caused by injections of kainate does not exactly mimic the damage observed in cerebral ischaemia. Nevertheless, given the possibility that administration of tPA to stroke patients might lead to increased protease levels in the brain parenchyma<sup>12</sup>, it seems prudent to examine the potential risk that tPA may cause neuronal death in the already vulnerable tissue.

**Stella E. Tsirka**

**Andrew D. Rogove**

**Sidney Strickland**

*Department of Pharmacology,  
University Medical Center at*

*Stony Brook,*

*Stony Brook,*

*New York 11794-8651, USA*

1. Sappino, A. et al. *J. Clin. Invest.* **92**, 679–685 (1993).
2. Qian, Z., Gilbert, M., Colicos, M., Kandel, E. & Kuhl, D. *Nature* **361**, 453–457 (1993).
3. Carroll, P., Tsirka, S., Richards, W., Frohman, M. & Strickland, S. *Development* **120**, 3173–3183 (1994).
4. Carmeliet, P. et al. *Nature* **368**, 419–424 (1994).
5. Tsirka, S., Gualandris, A., Amaral, D. & Strickland, S. *Nature* **377**, 340–344 (1995).
6. Andersson, P., Perry, V. & Gordon, S. *Neuroscience* **42**, 201–214 (1991).
7. Coyle, J., Molliver, M. & Kuhar, M. *J. Comp. Neurol.* **180**, 301–323 (1978).
8. Hekman, C. & Loskutoff, D. *Arch. Biochem. Biophys.* **262**, 199–210 (1988).
9. Lipton, S. & Rosenberg, P. *New Engl. J. Med.* **330**, 613–622 (1994).
10. Hacke, W. et al. *J. Am. Med. Assoc.* **274**, 1017–1025 (1995).
11. The NINDS rtPA Stroke Study Group *New Engl. J. Med.* **333**, 1581–1587 (1995).
12. Klatzo, I. *Basic Neurosci.* **1**, 313–322 (1975).




FIG. 1 Intra-hippocampal delivery of tPA restores susceptibility of hippocampal neurons to kainate injury in tPA<sup>-/-</sup> mice. Coronal sections through the hippocampus of tPA<sup>-/-</sup> mice stained with cresyl violet to reveal the status of neuronal cells. Mice were infused with PBS (top) or recombinant human tPA (bottom) for 2 days, injected unilaterally with kainate, and the infusion continued for 5 days further, at which point the mice were analysed. Percentage lengths of hippocampal subfields remaining intact after kainate infusion: 100% CA1, 91.2 ± 0.1% CA2/CA3 (top); 10.3 ± 4.4% CA1, 9.1 ± 5.2% CA2/CA3 (bottom). Arrowheads, site of injection; arrows, site of infusion. A micro-osmotic pump (Alzet Inc.) containing buffer (for control animals), or 100 µl human tPA (rtPA, 0.12 mg ml<sup>-1</sup>), was placed subcutaneously in the back of anaesthetized mice, and a brain infusion cannula connected to the pump was positioned at coordinates: bregma - 2.5 mm, medial-lateral - 0.5 mm and dorsoventral 1.6 mm, for delivery near the midline. The pump was allowed to infuse (0.5 µl h<sup>-1</sup>) for 2 days. Mice were then injected with kainate and analysed<sup>4</sup>.





FIG. 2 Plasminogen activator inhibitor type-1 prevents kainate-induced neuronal degeneration. Cresyl violet-stained coronal sections through the hippocampus of wild-type mice. The mice were infused with buffer (top), or mouse PAI-1 (0.12 mg ml<sup>-1</sup>, American Diagnostica; bottom) for 2 days, and kainate injected as in Fig. 1. Percentage lengths intact after kainate infusion: 6.3 ± 3.1% CA1, 18.1 ± 3.6% CA2/CA3 (top); 68.3 ± 7.4% CA1, 80.9 ± 4.1% CA2/CA3 (bottom). Arrowheads, site of injection; arrows, site of infusion.

**Correction**



In the Scientific Correspondence “Is ball lightning an electromagnetic knot?” by A. F. Rañada and J. L. Trueba (*Nature* **383**, 32; 1996), Fig. 1, a schematic version of the magnetic lines in the case  $n=1$ , was printed incorrectly. The correct version is shown above. In Fig. 2a, the origin on the ordinate axis should have read ‘1’.